




dNTP Mix dUTP - 10 mM Solution

Premix of 10 mM dATP, dCTP, dGTP and 20 mM dUTP

2'-Deoxyadenosine-5'-triphosphate, Sodium salt; 2'-Deoxycytidine-5'-triphosphate, Sodium salt; 2'-Deoxyguanosine-5'-triphosphate, Sodium salt; 2'-Deoxyuridine-5'-triphosphate, Sodium salt

Cat. No.	Amount
NU-1020S	200 µl
NU-1020L	1 ml



Structural formula of dNTP Mix dUTP - 10 mM Solution

For *in vitro* use only!

Shipping: shipped on blue ice

Storage Conditions: store at -20 °C

Short term exposure (up to 1 week cumulative) to ambient temperature possible. If stored as recommended, Jena Bioscience guarantees optimal performance of this product for 12 months after date of delivery.

Shelf Life: 12 months

Molecular Formula:

dATP: C₁₀H₁₆N₅O₁₂P₃ (free acid)

dCTP: C₉H₁₆N₃O₁₃P₃ (free acid)

dGTP: C₁₀H₁₆N₅O₁₃P₃ (free acid)

dUTP: C₉H₁₅N₂O₁₄P₃ (free acid)

Molecular Weight:

dATP: 491.18 g/mol (free acid)

dCTP: 467.15 g/mol (free acid)

dGTP: 507.18 g/mol (free acid)

dUTP: 468.14 g/mol (free acid)

Purity: ≥ 99 % (HPLC)

Form: clear aqueous solution

pH: 8.5 ± 0.2 (22 °C)

Description:

dNTP Mix incl. dUTP is a mixture of 10 mM ultrapure dATP, dCTP, and dGTP, and 20 mM dUTP supplied as clear aqueous solution (pH 8.5). dUTP can be used in place of dTTP in PCR and RT-PCR protocols to prevent carry-over contaminations from previous amplifications.

Selected References:

Erlich *et al.* (1988) Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* **29** (239):487.

Holland *et al.* (1991) Detection of specific polymerase chain reaction product by utilizing the 5'—3' exonuclease activity of *Thermus aquaticus* DNA polymerase. *Proc. Natl. Acad. Sci. USA* **88** (16):7276.

Sanger *et al.* (1977) DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463.